



New Approach to Fractional Milne-type Inequalities

Hüseyin Budak^{a,b,*}

^aDepartment of Mathematics, Faculty of Science and Arts, Kocaeli University, Kocaeli 41001, Türkiye.

^bDepartment of Mathematics, Saveetha School of Engineering, SIMATS, Saveetha University, Chennai 602105, Tamil Nadu, India.

Abstract

In the literature, there are several Milne-type inequalities by using the convex functions. In this study, we establish the upper and below bounds for fractional Milne-type inequalities by using functions whose second derivatives are bounded instead of convex functions. Moreover, we present new inequalities for Riemann integrals as special cases. We also present an example and a graph to illustrate the main results.

Keywords: Milne inequality, integral inequalities, bounded functions

2010 Mathematics Subject Classification: 26D07, 26D10, 26D15, 26A33

1. Introduction

The Hermite-Hadamard inequality, recognized as one of the foundational results concerning convex functions, possesses a clear geometric interpretation and has found numerous applications across various fields of mathematics. It has attracted significant attention within elementary mathematical analysis, prompting many researchers to explore generalizations, refinements, extensions, and analogous results for broader classes of functions, particularly through the lens of convexity.

The inequalities originally formulated by C. Hermite and J. Hadamard for convex functions hold a prominent place in the mathematical literature (see, for example, [12, p.137], [8]). These inequalities state that if $\varpi : I \rightarrow \mathbb{R}$ is a convex function on the interval I of real numbers and $a, b \in I$ with $a < b$, then

$$\varpi\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b \varpi(x) dx \leq \frac{\varpi(a) + \varpi(b)}{2}. \quad (1.1)$$

Both inequalities hold in the reversed direction if ϖ is concave. Some recent papers on Hermite-Hadamard inequalities, please refer to [1, 2, 14, 16]

In [6] and [7], Dragomir et al. proved the following results connected with the Hermite-Hadamard inequality:

*Corresponding author

Email address: hsyn.budak@gmail.com (Hüseyin Budak)

Theorem 1.1. Let $\varpi : [a, b] \rightarrow \mathbb{R}$ be a twice differentiable mapping such that there exists real constants m and M so that $m \leq \varpi'' \leq M$. Then, the following inequalities hold:

$$m \frac{(b-a)^2}{24} \leq \frac{1}{b-a} \int_a^b \varpi(\varkappa) d\varkappa - \varpi\left(\frac{a+b}{2}\right) \leq M \frac{(b-a)^2}{24}. \tag{1.2}$$

and

$$m \frac{(b-a)^2}{12} \leq \frac{\varpi(a) + \varpi(b)}{2} - \frac{1}{b-a} \int_a^b \varpi(\varkappa) d\varkappa \leq M \frac{(b-a)^2}{12} \tag{1.3}$$

In the following we will give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used further in this paper.

Definition 1.2. Let $\varpi \in L_1[a, b]$. The Riemann-Liouville integrals $J_{a+}^\alpha \varpi$ and $J_{b-}^\alpha \varpi$ of order $\alpha > 0$ with $\alpha \geq 0$ are defined by

$$J_{a+}^\alpha \varpi(\varkappa) = \frac{1}{\Gamma(\alpha)} \int_a^\varkappa (\varkappa - \varrho)^{\alpha-1} \varpi(\varrho) d\varrho, \quad \varkappa > a$$

and

$$J_{b-}^\alpha \varpi(\varkappa) = \frac{1}{\Gamma(\alpha)} \int_\varkappa^b (\varrho - \varkappa)^{\alpha-1} \varpi(\varrho) d\varrho, \quad \varkappa < b$$

respectively. Here, $\Gamma(\alpha)$ is the Gamma function and $J_{a+}^0 \varpi(\varkappa) = J_{b-}^0 \varpi(\varkappa) = \varpi(\varkappa)$.

For more information about fraction calculus please refer to [9, 10, 11, 13]

In [15], Sarikaya et al. first give the following interesting integral inequalities of Hermite-Hadamard type involving Riemann-Liouville fractional integrals.

Theorem 1.3. Let $\varpi : [a, b] \rightarrow \mathbb{R}$ be a convex function on $[a, b]$, then the following inequalities for fractional integrals hold:

$$\varpi\left(\frac{a+b}{2}\right) \leq \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a+}^\alpha \varpi(b) + J_{b-}^\alpha \varpi(a)] \leq \frac{\varpi(a) + \varpi(b)}{2} \tag{1.4}$$

with $\alpha > 0$.

Moreover, Dragomir give the following another version of Hermite-Hadamard inequality for Riemann-Liouville fractional integrals:

Theorem 1.4. [5] Let $\varpi : [a, b] \rightarrow \mathbb{R}$ be a convex function on $[a, b]$, then the following inequalities for fractional integrals hold:

$$\begin{aligned} \varpi\left(\frac{a+b}{2}\right) &\leq \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha} \left[J_{a+}^\alpha \varpi\left(\frac{a+b}{2}\right) + J_{b-}^\alpha \varpi\left(\frac{a+b}{2}\right) \right] \\ &\leq \frac{\varpi(a) + \varpi(b)}{2} \end{aligned} \tag{1.5}$$

Theorem 1.5. [3] Let $\varpi : [a, b] \rightarrow \mathbb{R}$ be a twice differentiable function with $a < b$ and $\varpi \in L_1[a, b]$. If ϖ'' is bounded, i.e. $m \leq \varpi''(\varrho) \leq M$, $\varrho \in (a, b)$, $m, M \in \mathbb{R}$, then we have the inequalities

$$\begin{aligned} &\frac{m(b-a)^2}{8(\alpha+2)} \\ &\leq \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha} \left[J_{a+}^\alpha \varpi\left(\frac{a+b}{2}\right) + J_{b-}^\alpha \varpi\left(\frac{a+b}{2}\right) \right] - \varpi\left(\frac{a+b}{2}\right) \\ &\leq \frac{M(b-a)^2}{8(\alpha+2)} \end{aligned}$$

and

$$\begin{aligned} & \frac{m(b-a)^2}{4(\alpha+2)} \\ & \leq \frac{\varpi(a) + \varpi(b)}{2} - \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[J_{a^+}^\alpha \varpi \left(\frac{a+b}{2} \right) + J_{b^-}^\alpha \varpi \left(\frac{a+b}{2} \right) \right] \\ & \leq \frac{M(b-a)^2}{4(\alpha+2)}. \end{aligned}$$

Now we present some Milne-type inequalities:

Theorem 1.6. [4] Let $\varpi : [a, b] \rightarrow \mathbb{R}$ be a differentiable mapping on (a, b) such that $\varpi \in L_1([a, b])$. If the function $|\varpi'|$ is convex on $[a, b]$, then we have the following Milne-type inequality

$$\begin{aligned} & \left| \frac{1}{3} \left[2\varpi(a) - \varpi \left(\frac{a+b}{2} \right) + 2\varpi(b) \right] - \frac{1}{b-a} \int_a^b \varpi(\varrho) d\varrho \right| \\ & \leq \frac{5(b-a)}{24} [|\varpi'(a)| + |\varpi'(b)|]. \end{aligned}$$

Theorem 1.7. [4] Assume that the assumptions of Theorem 1.6. Then we have

$$\begin{aligned} & \left| \frac{1}{3} \left[2\varpi(a) - \varpi \left(\frac{a+b}{2} \right) + 2\varpi(b) \right] \right. \\ & \quad \left. - \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[J_{a^+}^\alpha \varpi \left(\frac{a+b}{2} \right) + J_{b^-}^\alpha \varpi \left(\frac{a+b}{2} \right) \right] \right| \\ & \leq \frac{\alpha+4}{12(\alpha+1)} (b-a) [|\varpi'(a)| + |\varpi'(b)|], \end{aligned}$$

for $\alpha > 0$.

2. Main Results

In this section, we will prove an upper and below bound for the difference given in Theorem 1.7 by using the functions whose second derivatives are bounded.

Theorem 2.1. Let $\varpi : [a, b] \rightarrow \mathbb{R}$ be a twice differentiable function with $a < b$ and $\varpi \in L_1[a, b]$. If ϖ'' is bounded, i.e. $m \leq \varpi''(\varrho) \leq M$, $\varrho \in (a, b)$, $m, M \in \mathbb{R}$, then we have the inequalities

$$\begin{aligned} & \frac{\alpha+8}{8(\alpha+2)} (b-a)^2 m \tag{2.1} \\ & \leq \frac{1}{3} \left[2\varpi(a) - \varpi \left(\frac{a+b}{2} \right) + 2\varpi(b) \right] \\ & \quad - \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[J_{a^+}^\alpha \varpi \left(\frac{a+b}{2} \right) + J_{b^-}^\alpha \varpi \left(\frac{a+b}{2} \right) \right] \\ & \leq \frac{\alpha+8}{8(\alpha+2)} (b-a)^2 M. \end{aligned}$$

Proof. From the definition of Riemann-Liouville fractional integrals, we get

$$\begin{aligned} & \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[J_{a+}^\alpha \varpi \left(\frac{a+b}{2} \right) + J_{b-}^\alpha \varpi \left(\frac{a+b}{2} \right) \right] \\ &= \frac{2^{\alpha-1}\alpha}{(b-a)^\alpha} \left[\int_a^{\frac{a+b}{2}} \left(\frac{a+b}{2} - \varkappa \right)^{\alpha-1} \varpi(\varkappa) d\varkappa + \int_{\frac{a+b}{2}}^b \left(\varkappa - \frac{a+b}{2} \right)^{\alpha-1} \varpi(\varkappa) d\varkappa \right] \\ &= \frac{2^{\alpha-1}\alpha}{(b-a)^\alpha} \int_a^{\frac{a+b}{2}} [\varpi(\varkappa) + \varpi(a+b-\varkappa)] \left(\frac{a+b}{2} - \varkappa \right)^{\alpha-1} d\varkappa. \end{aligned} \tag{2.2}$$

Using the identity (2.2), we get

$$\begin{aligned} & \frac{1}{3} \left[2\varpi(a) - \varpi \left(\frac{a+b}{2} \right) + 2\varpi(b) \right] \\ & - \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[J_{a+}^\alpha \varpi \left(\frac{a+b}{2} \right) + J_{b-}^\alpha \varpi \left(\frac{a+b}{2} \right) \right] \\ &= \frac{1}{3} \left[2\varpi(a) - \varpi \left(\frac{a+b}{2} \right) + 2\varpi(b) \right] \\ & - \frac{2^{\alpha-1}\alpha}{(b-a)^\alpha} \int_a^{\frac{a+b}{2}} [\varpi(\varkappa) + \varpi(a+b-\varkappa)] \left(\frac{a+b}{2} - \varkappa \right)^{\alpha-1} d\varkappa \\ &= \frac{2^{\alpha-1}\alpha}{3(b-a)^\alpha} \int_a^{\frac{a+b}{2}} \left[\begin{aligned} & 4\varpi(a) - 2\varpi \left(\frac{a+b}{2} \right) + 4\varpi(b) \\ & - 3\varpi(\varkappa) - 3\varpi(a+b-\varkappa) \end{aligned} \right] \left(\frac{a+b}{2} - \varkappa \right)^{\alpha-1} d\varkappa. \end{aligned} \tag{2.3}$$

Using the facts that

$$\varpi(\varkappa) - \varpi(a) = \int_a^\varkappa \varpi'(\varrho) d\varrho$$

and

$$\varpi(b) - \varpi(a+b-\varkappa) = \int_{a+b-\varkappa}^b \varpi'(\varrho) d\varrho,$$

we get

$$\begin{aligned} & \varpi(a) + \varpi(b) - \varpi(\varkappa) - \varpi(a+b-\varkappa) \\ &= \int_{a+b-\varkappa}^b \varpi'(\varrho) d\varrho - \int_a^\varkappa \varpi'(\varrho) d\varrho = \int_a^\varkappa \varpi'(a+b-u) du - \int_a^\varkappa \varpi'(\varrho) d\varrho = \int_a^\varkappa [\varpi'(a+b-\varrho) - \varpi'(\varrho)] d\varrho. \end{aligned} \tag{2.4}$$

On the other hand, we have

$$\varpi(a) - \varpi \left(\frac{a+b}{2} \right) = - \int_a^{\frac{a+b}{2}} \varpi'(\varrho) d\varrho \tag{2.5}$$

and

$$\varpi(b) - \varpi\left(\frac{a+b}{2}\right) = \int_{\frac{a+b}{2}}^b \varpi'(\varrho) d\varrho. \tag{2.6}$$

By (2.5) and (2.6), we get

$$\begin{aligned} & \varpi(a) + \varpi(b) - 2\varpi\left(\frac{a+b}{2}\right) \\ &= \int_{\frac{a+b}{2}}^b \varpi'(\varrho) d\varrho - \int_a^{\frac{a+b}{2}} \varpi'(\varrho) d\varrho = \int_a^{\frac{a+b}{2}} \varpi'(a+b-\varrho) d\varrho - \int_a^{\frac{a+b}{2}} \varpi'(\varrho) d\varrho = \int_a^{\frac{a+b}{2}} [\varpi'(a+b-\varrho) - \varpi'(\varrho)] d\varrho. \end{aligned} \tag{2.7}$$

We also have

$$\varpi'(a+b-\varrho) - \varpi'(\varrho) = \int_{\varrho}^{a+b-\varrho} \varpi''(u) du. \tag{2.8}$$

By using the equality (2.8) and $m < \varpi''(u) < M, u \in (a, b)$, we obtain,

$$m \int_{\varrho}^{a+b-\varrho} du \leq \int_{\varrho}^{a+b-\varrho} \varpi''(u) du \leq M \int_{\varrho}^{a+b-\varrho} du$$

i.e.

$$m(a+b-2\varrho) \leq \varpi'(a+b-\varrho) - \varpi'(\varrho) \leq M(a+b-2\varrho). \tag{2.9}$$

By integrating the inequality (2.9) with respect to ϱ on $[a, \varkappa]$, from (2.4), we get

$$\begin{aligned} & m \left[\left(\frac{b-a}{2}\right)^2 - \left(\frac{a+b}{2} - \varkappa\right)^2 \right] \\ & \leq \varpi(a) + \varpi(b) - \varpi(\varkappa) - \varpi(a+b-\varkappa) \\ & \leq M \left[\left(\frac{b-a}{2}\right)^2 - \left(\frac{a+b}{2} - \varkappa\right)^2 \right]. \end{aligned} \tag{2.10}$$

That is,

$$\begin{aligned} & m(\varkappa - a)(b - \varkappa) \\ & \leq \varpi(a) + \varpi(b) - \varpi(\varkappa) - \varpi(a+b-\varkappa) \\ & \leq M(\varkappa - a)(b - \varkappa). \end{aligned} \tag{2.11}$$

Similarly, integrating the inequality (2.9) with respect to ϱ on $[a, \frac{a+b}{2}]$, from (2.7), we obtain

$$m \left(\frac{b-a}{2}\right)^2 \leq \varpi(a) + \varpi(b) - 2\varpi\left(\frac{a+b}{2}\right) \leq M \left(\frac{b-a}{2}\right)^2. \tag{2.12}$$

By using the inequalities (2.11) and (2.12), we get

$$\begin{aligned}
 & 3m(\varkappa - a)(b - \varkappa) + m\left(\frac{b - a}{2}\right)^2 \tag{2.13} \\
 & \leq 4\varpi(a) - 2\varpi\left(\frac{a + b}{2}\right) + 4\varpi(b) - 3\varpi(\varkappa) - 3\varpi(a + b - \varkappa) \\
 & \leq 3M(\varkappa - a)(b - \varkappa) + M\left(\frac{b - a}{2}\right)^2.
 \end{aligned}$$

Multiplying the inequality (2.13) by $\frac{2^{\alpha-1}\alpha}{(b-a)^\alpha} \left(\frac{a+b}{2} - \varkappa\right)^{\alpha-1}$ and integrating the resultant inequality with respect to \varkappa on $\left[a, \frac{a+b}{2}\right]$, we establish

$$\begin{aligned}
 & \frac{m2^{\alpha-1}\alpha}{(b-a)^\alpha} \int_a^{\frac{a+b}{2}} \left[3(\varkappa - a)(b - \varkappa) + \left(\frac{b - a}{2}\right)^2 \right] \left(\frac{a + b}{2} - \varkappa\right)^{\alpha-1} d\varkappa \\
 & \leq \frac{2^{\alpha-1}\alpha}{3(b-a)^\alpha} \int_a^{\frac{a+b}{2}} \left[4\varpi(a) - 2\varpi\left(\frac{a+b}{2}\right) + 4\varpi(b) - 3\varpi(\varkappa) - 3\varpi(a + b - \varkappa) \right] \left(\frac{a + b}{2} - \varkappa\right)^{\alpha-1} d\varkappa \\
 & \leq \frac{M2^{\alpha-1}\alpha}{(b-a)^\alpha} \int_a^{\frac{a+b}{2}} \left[3(\varkappa - a)(b - \varkappa) + \left(\frac{b - a}{2}\right)^2 \right] \left(\frac{a + b}{2} - \varkappa\right)^{\alpha-1} d\varkappa.
 \end{aligned}$$

By the equality (2.3) and equality

$$\begin{aligned}
 & \frac{2^{\alpha-1}\alpha}{(b-a)^\alpha} \int_a^{\frac{a+b}{2}} \left[3(\varkappa - a)(b - \varkappa) + \left(\frac{b - a}{2}\right)^2 \right] \left(\frac{a + b}{2} - \varkappa\right)^{\alpha-1} d\varkappa \\
 & = \frac{\alpha + 8}{8(\alpha + 2)} (b - a)^2,
 \end{aligned}$$

we get the desired inequality (2.1).

This completes the proof. □

Remark 2.2. If we choose $\alpha = 1$ in Theorem 2.1, then we have the following inequality

$$\begin{aligned}
 & \frac{3}{8} (b - a)^2 m \\
 & \leq \frac{1}{3} \left[2\varpi(a) - \varpi\left(\frac{a + b}{2}\right) + 2\varpi(b) \right] - \frac{1}{b - a} \int_a^b \varpi(\varrho) d\varrho \\
 & \leq \frac{3}{8} (b - a)^2 M.
 \end{aligned}$$

This is a new contribution for the literature. This Milne inequality holds for the functions whose second derivatives are bounded.

Example 2.3. Let consider the function $\varpi : [-1, 2] \rightarrow \mathbb{R}$ defined by $\varpi(\varrho) = \varrho^5$. It is clear that $\varpi''(\varrho) = 20\varrho^3$ which is not convex on $[-1, 2]$ but bounded such that $-20 \leq \varpi''(\varrho) \leq 160$. Therefore we have the left hand side and right hand side of the inequality (2.1) as

$$\frac{\alpha + 8}{8(\alpha + 2)} (b - a)^2 m = -\frac{45(\alpha + 8)}{2(\alpha + 2)}$$

and

$$\frac{\alpha + 8}{8(\alpha + 2)} (b - a)^2 M = \frac{180(\alpha + 8)}{(\alpha + 2)}$$

respectively. We also have

$$\frac{1}{3} \left[2\varpi(a) - \varpi\left(\frac{a+b}{2}\right) + 2\varpi(b) \right] = \frac{1}{3} \left[62 - \frac{1}{32} \right] = \frac{661}{32}.$$

By definition of Riemann-Liouville fractional integrals, we have

$$\begin{aligned} J_{a+}^{\alpha} \varpi\left(\frac{a+b}{2}\right) &= \frac{1}{\Gamma(\alpha)} \int_{-1}^{\frac{1}{2}} \left(\frac{1}{2} - \varrho\right)^{\alpha-1} \varrho^5 d\varrho \\ &= \frac{1}{2^{\alpha+5}\Gamma(\alpha)} \left[\frac{3^{\alpha}}{\alpha} - \frac{5 \cdot 3^{\alpha+1}}{\alpha+1} + \frac{10 \cdot 3^{\alpha+2}}{\alpha+2} - \frac{10 \cdot 3^{\alpha+3}}{\alpha+3} + \frac{5 \cdot 3^{\alpha+4}}{\alpha+4} - \frac{3^{\alpha+5}}{\alpha+5} \right] \end{aligned}$$

and

$$\begin{aligned} J_{b-}^{\alpha} \varpi\left(\frac{a+b}{2}\right) &= \frac{1}{\Gamma(\alpha)} \int_{\frac{1}{2}}^2 \left(\varrho - \frac{1}{2}\right)^{\alpha-1} \varrho^5 d\varrho \\ &= \frac{1}{2^{\alpha+5}\Gamma(\alpha)} \left[\frac{3^{\alpha}}{\alpha} + \frac{5 \cdot 3^{\alpha+1}}{\alpha+1} + \frac{10 \cdot 3^{\alpha+2}}{\alpha+2} + \frac{10 \cdot 3^{\alpha+3}}{\alpha+3} + \frac{5 \cdot 3^{\alpha+4}}{\alpha+4} + \frac{3^{\alpha+5}}{\alpha+5} \right]. \end{aligned}$$

Thus, the midterm of the inequality (2.1) becomes

$$\begin{aligned} &\frac{1}{3} \left[2\varpi(a) - \varpi\left(\frac{a+b}{2}\right) + 2\varpi(b) \right] - \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^{\alpha}} \left[J_{a+} \varpi\left(\frac{a+b}{2}\right) + J_{b-} \varpi\left(\frac{a+b}{2}\right) \right] \\ &= \frac{661}{32} - \frac{\alpha}{32} \left[\frac{1}{\alpha} + \frac{90}{\alpha+2} + \frac{405}{\alpha+4} \right]. \end{aligned}$$

From the inequality (2.1), we have

$$-\frac{45(\alpha + 8)}{2(\alpha + 2)} \leq \frac{661}{32} - \frac{\alpha}{32} \left[\frac{1}{\alpha} + \frac{90}{\alpha + 2} + \frac{405}{\alpha + 4} \right] \leq \frac{180(\alpha + 8)}{(\alpha + 2)}. \tag{2.14}$$

One can see the validity of the inequality (2.14) in Figure 1.




Figure 1: Comparison of the terms of the inequality (2.14)

3. Concluding Remarks

In this study, we obtained fractional Milne-type inequalities by using the functions whose second derivatives are bounded instead of convex functions. The method in this paper may applied for the any other type fractional integrals in the future works

References

- [1] N. Azzouz and B. Benaissa. Exploring hermite–hadamard-type inequalities via ψ -conformable fractional integral operators. *Journal of Inequalities and Mathematical Analysis*, 1(1):15–27, 2025. 1
- [2] B. Benaissa and N. Azzouz. Hermite-hadamard-fejér type inequalities for h-convex functions involving ψ -hilfer operators. *Journal of Inequalities and Mathematical Analysis*, 1(2):113–123, 2025. 1
- [3] H. Budak, H. Kara, M. Sarikaya, and M. Kiriş. New extensions of the hermite-hadamard inequalities involving riemann-liouville fractional integrals. *Miskolc Mathematical Notes*, 21(2):665–678, 2020. 1.5
- [4] H. Budak, P. Kösem, and H. Kara. On new milne-type inequalities for fractional integrals. *Journal of inequalities and applications*, 2023(1):10, 2023. 1.6, 1.7
- [5] S. S. Dragomir. Some inequalities of hermite-hadamard type for symmetrized convex functions and riemann-liouville fractional integrals. *RGMA Res. Rep. Coll*, 20:15, 2017. 1.4
- [6] S. S. Dragomir, P. Cerone, and A. Sofo. Some remarks on the midpoint rule in numerical integration. *Studia Univ. Babeş-Bolyai, Math.*, XLV(1):63–74, 2000. 1
- [7] S. S. Dragomir, P. Cerone, and A. Sofo. Some remarks on the trapezoid rule in numerical integration. *Indian J. Pure Appl. Math.*, 31(5):475–494, 2000. 1
- [8] S. S. Dragomir and C. Pearce. Selected topics on hermite-hadamard inequalities and applications. *Science direct working paper*, (S1574-0358):04, 2003. 1
- [9] R. Gorenflo and F. Mainardi. Fractional calculus: integral and differential equations of fractional order. In *Fractals and fractional calculus in continuum mechanics*, pages 223–276. Springer, 1997. 1
- [10] A. Kilbas. *Theory and applications of fractional differential equations*, volume 204. Elsevier, 2006. 1
- [11] K. S. Miller and B. Ross. *An introduction to the fractional calculus and fractional differential equations*. John Wiley & Sons, 1993. 1
- [12] J. E. Pečarić and Y. L. Tong. *Convex functions, partial orderings, and statistical applications*. Academic press, 1992. 1
- [13] I. Podlubny. *Fractional differential equations: an introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications*, volume 198. elsevier, 1998. 1
- [14] M. Z. Sarikaya. Generalized hermite-hadamard integral on intervals. *Turkish Journal of Inequalities*, 9(2):70–78, 2025. 1
- [15] M. Z. Sarikaya, E. Set, H. Yaldiz, and N. Başak. Hermite–hadamard’s inequalities for fractional integrals and related fractional inequalities. *Mathematical and computer modelling*, 57(9-10):2403–2407, 2013. 1
- [16] Ö. Ustaoglu, H. Kara, and H. Budak. Hermite-hadamard inequalities involving fractional conformable integral operators for the product of two interval-valued lr-convex functions. *Communications in Advanced Mathematical Sciences*, 8(3):136–150, 2025. 1